

**Neomycin Sulfate and Oxytetracycline Calf Supplement II
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated calf feed for use in calf milk replacers or starter feeds.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....10 grams/lb.
Oxytetracycline hydrochloride.....10 grams/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....XX.XX%
NPN, (Max)¹..... X.XX%
Crude Fat, (Min)..... X.XX%
Crude Fiber, (Max)..... X.XX%
Calcium, (Min)..... X.XX%
Calcium, (Max)..... X.XX%
Phosphorus, (Min)..... X.XX%
Salt, (Min)¹..... X.XX%
Salt, (Max)¹..... X.XX%
Sodium, (Min)²..... X.XX%
Sodium, (Max)²..... X.XX%
Potassium, (Min)¹..... X.XX%
Vitamin A, (Min)³..... ____IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. For the manufacture of Type C medicated feeds of less than 100g/ton neomycin and oxytetracycline from this Type B feed, it is recommended that the Type B feed be further diluted before mixing the Type C feed. An example of further dilution would be a ratio of 1:10 of Type B medicated feed:unmedicated feed.

Indications for Use	Oxytetracycline and Neomycin Sulfate Amount	Mixing Directions
Calves (up to 250 lbs): For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin	10 mg/lb of body weight daily	Mix an amount of Type B feed in a Type C medicated feed, as starter or milk replacer, to provide 10 mg/lb body weight daily*
<p align="center">Residue Warning: Withdraw 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residue. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.</p>		

*For example, if an animal weighs 200 lb, consuming 4.00 lb of Type C feed daily, the feed should contain 1000 g/ton oxytetracycline and neomycin sulfate. For 1000 g/ton, mix 100 lb of Type B feed per ton of Type C feed.

Net Weight _____lb. (_____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000