

**Neomycin Sulfate and Oxytetracycline Turkey Ration C
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)
drug to use by or on the order of a licensed veterinarian.**

INDICATIONS FOR USE

For control of infectious synovitis caused by Mycoplasma synoviae susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....200 grams/ton.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 200 grams oxytetracycline hydrochloride/ton.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously for 7-14 days as the sole ration.

RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.
Withdraw 5 days before slaughter.

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000