

**Neomycin Sulfate and Oxytetracycline Turkey Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated turkey feed.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....10 grams/lb.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 10 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below.

Indications for Use	Oxytetracycline and Neomycin Sulfate Amount	Mixing Directions
For control of hexamitiasis caused by <u>Hexamita meleagridis</u> susceptible to oxytetracycline.	100 g/ton of neomycin and oxytetracycline	Mix 10 lb of this Type B medicated feed with 1990 lb of non-medicated feed to make one ton of Type C medicated feed containing 100 g/ton of neomycin and oxytetracycline
For control of infectious synovitis caused by <u>Mycoplasma synoviae</u> susceptible to oxytetracycline.	200 g/ton of neomycin and oxytetracycline	Mix 20 lb of this Type B medicated feed with 1980 lb of non-medicated feed to make one ton of Type C medicated feed containing 200 g/ton of

		neomycin and oxytetracycline
For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.	25mg/ lb body weight/day	Mix 100 lb of this Type B medicated feed with 1900 lb of non-medicated feed to make one ton of Type C medicated feed containing 1000 g/ton of neomycin and oxytetracycline*
 Residue Warning: Do not feed to turkeys producing eggs for human consumption. At 200 g/ton use level or higher, withdraw 5 days before slaughter. Zero-day withdrawal period at 100 g/ton use level. 		

*A 20lb animal consuming 1 lb of this Type C feed will receive the required amount of oxytetracycline and neomycin sulfate.

Net Weight _____lb. (_____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000