

**Chlortetracycline Turkey Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated turkey feed.

ACTIVE DRUG INGREDIENT

Chlortetracycline.....2-40 g/lb*

GUARANTEED ANALYSIS

Crude Protein, not less than.....%
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Lysine (min).....%
Methionine (min).....%
Calcium (min).....%
Calcium (max).....%
Phosphorus(min).....%
Salt¹ (min).....%
Salt¹ (max).....%
Sodium² (min).....%
Sodium² (max).....%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Blend with grains and non-medicated feed ingredients to provide a complete feed with the following grams of chlortetracycline per ton of complete feed.

Indications for Use

**Use Levels of
Chlortetracycline**

TURKEYS

For Turkeys: Mix 100 pounds of Type B (2 g/lb) or 5 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for the control of infectious synovitis caused by *Mycoplasma synoviae*, susceptible to chlortetracycline containing 200 g/ton to be fed continuously for 7 to 14 days.

200 g/ton

WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.

For Turkeys: Mix 200 pounds of Type B (2 g/lb) or 10 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for the control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline containing 400 g/ton to be fed continuously for 7 to 14 days.

400 g/ton

WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.

For Turkeys: Turkeys poults not over 4 weeks of age: Mix 200 pounds of Type B (2 g/lb) or 10 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for the reduction of mortality due to paratyphoid caused by Salmonella typhimurium susceptible to chlortetracycline containing 400 g/ton.

400 g/ton

WARNING: Zero-day withdrawal period.

For Turkeys: Mix an amount of Type B feed to provide 25 mg/lb body weight/day varying with body weight and feed consumption with non-medicated ingredients to make a ton of turkey feed in the resulting Type C medicated feed for the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline to be fed continuously for 7-14 days.

25 mg/lb body weight/day

WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.

*The final printed label must bear a single drug concentration.

Net Weight _____ lb (_____ kg)
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000