

DERA/BOV COCCI — CATTLE FEED
TYPE B Medicated Feed
(chlortetracycline and lasalocid Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

ACTIVE DRUG INGREDIENTS*

Chlortetracycline^a 4000 to 80000 g/ton**
Lasalocid^b (as lasalocid sodium)..... 181.8 to 3636.4 g/ton**

GUARANTEED ANALYSIS

Crude protein, not less than %
(This includes not more than _____ % equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt², not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 4000 grams chlortetracycline per ton and 30 to 181.8 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Type B Concentration, g/ton		Type B per ton of Type C, lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
				Chlortetracycline	Lasalocid
CTC	Lasalocid				
66667	3030	120	1880	4000	181.8
66667	3030	60	1940	2000	90.9
66667	3030	30	1970	1000	45.5
66667	3030	20	1980	667	30.3
40000	1818	200	1800	4000	181.8
40000	1818	100	1900	2000	90.9
40000	1818	50	1950	1000	45.5
40000	1818	33.3	1966.7	667	30.3

Caution

Safety of lasalocid for use in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type B medicated feed must bear a single concentration for each drug.

**The ratio between chlortetracycline and lasalocid must be 22:1.

^aDeracin® is the proprietary name of chlortetracycline Type A medicated article (ANADA 200-510).

^bBOVATEC is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Pharmgate LLC [March 2017]