

BLUE BIRD LABELING
chlortetracycline hydrochloride
Type B Medicated Turkey Feed – Bluecomb

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline in turkeys.

Active Drug Ingredient

Chlortetracycline* 5,001 – 80,000 g/ton**

Guaranteed Analysis

Crude Protein (Min) _____ %
 Lysine (Min) _____ %
 Methionine (Min) _____ %
 Crude Fat (Min) _____ %
 Crude Fiber (Max) _____ %
 Calcium (Min) _____ %
 Calcium (Max) _____ %
 Phosphorus (Min) _____ %
 Salt^a (Min) _____ %
 Salt^a (Max) _____ %
 Sodium^b (Min) _____ %
 Sodium^b (Max) _____ %

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	200	1800	1000
20,000 (10 g/lb)	100	1900	1000
40,000 (20 g/lb)	50	1950	1000
10,000 (5 g/lb)	1000	1000	5000
20,000 (10 g/lb)	500	1500	5000
40,000 (20 g/lb)	250	1750	5000

The resulting Type C medicated feed should be fed to turkeys for 7 to 14 days, to provide chlortetracycline at the rate of 25 mg per pound of body weight per day.

Warning

Do not feed to turkeys producing eggs for human consumption.

Manufactured By:

Blue Bird Feed Mill
 City, State, Zip

NET WEIGHT ON BAG OR BULK

*Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).

**The final printed feed label should list only a single drug concentration.