

BLUE BIRD LABELING
chlortetracycline hydrochloride
Type C Medicated Psittacine Bird Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by *Chlamydia psittaci* sensitive to chlortetracycline.

Active Drug Ingredient

Chlortetracycline* 10 mg/g (1.0%)

Guaranteed Analysis

Crude Protein (Min) _____%

Crude Fat (Min)..... _____%

Crude Fiber (Max) _____%

Moisture (Max)..... _____%

Ingredients

Ingredients as defined by AAFCO.

Feeding Directions

Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.

Warning

Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials.

Caution

Aspergilliosis may occur following prolonged treatment.

Manufactured By:

Blue Bird Feed Mill

City, State, Zip

NET WEIGHT ON BAG OR BULK

*Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).