

Chlortetracycline and Sulfamethazine Cattle Feed
Type B Medicated Cattle Feed
(chlortetracycline and sulfamethazine Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in the maintenance of weight gains in the presence of respiratory disease such as shipping fever.

Active Drug Ingredient

Chlortetracycline^a.....3,000 - 20,000 g/ton (1.5 to 10 g/lb)*
Sulfamethazine^a.....3,000 - 20,000 g/ton (1.5 to 10 g/lb)*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
NPN ¹ , not more than	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than	_____	%
Calcium ¹ , not less than.....	_____	%
Calcium ¹ , not more than.....	_____	%
Phosphorus ¹ , not less than.....	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Potassium ¹ , not less than.....	_____	%
Vitamin A ¹ , not less than	_____	IU/lb

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated cattle feed.

The following table provides examples of mixing rates:

Type B CTC / SMZ concentration (g/lb)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC / SMZ concentration (g/ton)	Amount of Type C feed per head per day (lbs)
2 / 2	17.	1982.5	35 / 35	20
2 / 2	87.	1912.5	175 / 175	4
2 / 2	700	1300	1400 / 1400	0.5
3.5 / 3.5	10	1990	35 / 35	20
3.5 / 3.5	50	1950	175 / 175	4
3.5 / 3.5	400	1600	1400 / 1400	0.5
10 / 10	3.5	1996.5	35 / 35	20
10 / 10	17.	1982.5	175 / 175	4
10 / 10	140	1860	1400 / 1400	0.5

The resulting Type C medicated feed will furnish 350 mg chlortetracycline and 350 mg sulfamethazine per head per day and should be fed for 28 days.

Warning

Withdraw 7 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Label must show actual concentration of chlortetracycline (CTC) and sulfamethazine (SMZ), which will vary with feed consumption to provide 350 mg of each drug/head/day.

^a Pennchlor S 70® is the proprietary name of chlortetracycline and sulfamethazine Type A medicated article.

version 2/17/16