

Product Profile

Deracin® 90

Chlortetracycline Type A Medicated Article

Product Description

- **Antibacterial premix for oral administration in feed to cattle, swine, chickens, turkeys, ducks, and sheep**
- **Generic equivalent (ANADA 200-510) to Aureomycin®**
- **Each pound of premix contains 90 grams of chlortetracycline activity**



Formulation

- Free-flowing feed meal medication.
- **Ingredients:** Chlortetracycline calcium complex, dried *Streptomyces aureofaciens* fermentation product in a carrier suitable for incorporation in feed.

FDA Status

- **CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.
- Type A Medicated Article for use in the manufacture of medicated dry feeds (not for use in liquid feeds).
- Category I drug; does not require a feedmill license.

Precautions

- 0-days (no withdrawal needed).
- A withdrawal period has not been established for this product in preruminating calves.
- Do not use in calves to be processed for veal.
- Do not feed to ducks or turkeys producing eggs for human consumption.

Indications (Partial List, See Back)

- **Cattle:** For the reduction of the incidence of liver abscesses; treatment and control of bacterial pneumonia (*Pasteurella* spp.); treatment of bacterial enteritis (*E. coli*);- control of active infection of anaplasmosis (*Anaplasma marginale*).
- **Swine:** Treatment of bacterial enteritis (*E. coli*, *Salmonella choleraesuis*) and bacterial pneumonia (*Pasteurella multocida*); control of ileitis (*Lawsonia intracellularis*); control of leptospirosis (*Leptospira pomona*); reducing the incidence of jowl abscesses (Group E streptococci).
- **Chickens:** Control of infectious synovitis (*M. synoviae*); control of chronic respiratory disease (CRD) and air sac infection (*Mycoplasma gallisepticum*, *E. coli*); reduction of mortality due to *E. coli*.
- **Turkeys:** Control of infectious synovitis (*M. synoviae*); control of hexamitiasis (*Hexamita meleagrides*); reduction of mortality due to paratyphoid (*Salmonella typhimurium*); control of complicating bacteria associated with bluecomb (transmissible enteritis, coronaviral enteritis).

Key Features

- **Broad spectrum, effective against both Gram-positive and Gram-negative organisms that can cause respiratory, enteric, or reproductive diseases.**
- **Readily absorbed, providing effective blood and lung tissue concentrations.**
- **Cattle industry product standard for control of anaplasmosis.**
- **Can be top dressed in cattle feeds.**
- **Can be fed to cattle on pasture with blue bird free choice mineral formulation.**
- **Versatile, with wide range of approved combinations including Deccox® and Bovatec®.**
- **Convenient, cost-effective treatment option vs injectable therapy.**
- **Wide safety margin.**
- **High-quality manufacturing at US Pharmgate facilities according to FDA requirements.**

The labeling contains complete use information, including any cautions and warnings. Always read, understand and follow the labeling and use directions. See the reverse side for use directions and additional information.



Pharmgate
ANIMAL HEALTH

1800 Sir Tyler Drive
Wilmington, NC | 28405
910.679.8364
Pharmgate.com

Deracin® 90 Meal

Chlortetracycline Type A Medicated Article

Active Drug Ingredients:		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 90 grams Chlortetracycline Hydrochloride/lb.														
Ingredients:		Dried <i>Streptomyces aureofaciens</i> fermentation product in a carrier suitable for incorporation in feed.														
For Use in the Manufacture of Medicated Feeds																
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS																
Use directions:		Mix sufficient Deracin® 90 Meal Type A Medicated Article to supply desired concentration of chlortetracycline per ton with part of the feed ingredients to make a preblend. Add the remainder of the ingredients and mix thoroughly. For specific use levels, see Indications .														
Mixing directions:		<table border="1"> <thead> <tr> <th>Level desired grams per ton</th> <th>Amount of medicated article per ton*</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>9 oz</td> </tr> <tr> <td>100</td> <td>1 lb 2 oz</td> </tr> <tr> <td>200</td> <td>2 lb 4 oz</td> </tr> <tr> <td>400</td> <td>4 lb 8 oz</td> </tr> <tr> <td>500</td> <td>5 lb 9 oz</td> </tr> </tbody> </table>	Level desired grams per ton	Amount of medicated article per ton*	50	9 oz	100	1 lb 2 oz	200	2 lb 4 oz	400	4 lb 8 oz	500	5 lb 9 oz	*It is recommended that 1 pound 2 ounces of Deracin® 90 Meal Type A Medicated Article be diluted with 2 pounds 14 ounces of one of the feed ingredients to form a 4 pound working premix. Use 2 pounds of the working premix to make a preblend (see Use Directions) for a Type C feed containing 50g chlortetracycline / ton of feed	
Level desired grams per ton	Amount of medicated article per ton*															
50	9 oz															
100	1 lb 2 oz															
200	2 lb 4 oz															
400	4 lb 8 oz															
500	5 lb 9 oz															
Indications For Use	Chlortetracycline mg per lb body weight per day	Indications For Use	In complete feed chlortetracycline g per ton													
Cattle Beef Cattle (over 700lb): control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. Beef and Non-Lactating Dairy Cattle: As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline when delivered in a free-choice feed. Free-choice feed must be manufactured under a medicated feed mill license utilizing an FDA approved formulation. Calves, Beef, and Non-Lactating Dairy Cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Feed for not more than 5 days. The appropriate amount of Deracin® containing feed supplement may be mixed in the cattle's daily ration or administered as a top dress. If the Deracin® containing feed supplement is administered as a top dress, it must be spread uniformly on top of the ration and sufficient space must be provided so that all cattle can eat at the same time.	0.5 0.5-2.0 10	Swine Reduction in the incidence of cervical lymphadenitis (fowl abscesses) caused by Group E. <i>Streptococci</i> susceptible to chlortetracycline. Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. Feed continuously for not more than 14 days. Ducks Control and treatment of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending upon age and severity of disease. Feed for not more than 21 days. Chickens Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. Feed for 5 days.	50-100 400 200-400 100-200 200-400 500													
Swine Control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Note: This drug level is equivalent to approximately 400 grams per ton, depending on feed consumption and body weight). Feed for not more than 14 days.	10	Turkeys Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	200 400 400													
Turkeys Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	25															
Indications For Use	mg per head per day	Indications For Use	mg per g feed													
Cattle Growing Cattle (over 400 lb): For the reduction of the incidence of liver abscesses. Beef Cattle: Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline. Beef Cattle (under 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	70 350 350	Psittacine birds Warning: Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials. Caution: Aspergillosis may occur following prolonged treatment. Treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.	10													
Sheep Breeding Sheep: Reduction in the incidence of (vibriotic) abortions caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.	80															
Warning: A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.																
Store below 25°C (77°F), excursions permitted to 40°C (104°F). Tightly reseal opened bags.																
ANADA 200-510, Approved by FDA																

Rev. 10-16
Bag D90

