

bacitracin methylenedisalicylate

Type C medicated feed for beef steers & heifers fed in confinement for slaughter – Liver Abscess 250
(bacitracin Type C medicated feed)

For reduction in the number of liver condemnations due to abscesses in beef steers and heifers fed in confinement for slaughter.

Active Drug Ingredient

Bacitracin (as feed grade bacitracin methylenedisalicylate)* g/ton**

Guaranteed Analysis

Crude Protein (Min) %
 NPN^a (Max) %
 Crude Fat (Min) %
 Crude Fiber (Max) %
 Calcium^a (Min) %
 Calcium^a (Max) %
 Phosphorus^a (Min) %
 Salt^a (Min) %
 Salt^a (Max) %
 Sodium^b (Min) %
 Sodium^b (Max) %
 Potassium^a (Min) %
 Vitamin A^a IU/lb

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Feed ingredients, as defined by AAFCO, must be listed on the final printed Type C medicated feed label by their common or usual names in descending order or predominance by weight.

Feeding Directions

Administer continuously for 5 days to provide 250 mg bacitracin per head per day, then discontinue for subsequent 25 days; repeat the pattern during the feeding period. ___^{***} pounds of this feed will medicate one animal.

Manufactured By:

Blue Bird Feed Mill
City, State, Zip

NET WEIGHT ON BAG OR BULK

Lot # _____

*Pennitracin MD is the proprietary name of bacitracin methylenedisalicylate formulation (Type A Medicated Article).

**Label must show actual concentration of bacitracin, which will vary with consumption to provide 250 mg/head/day.

***The following table provides examples of feeding rates and amount of drug per ton of feed. Third column represents number of pounds of feed that will medicate one animal.

Drug level in feed (g/ton)	mg bacitracin/head/day	*** Pounds of feed/head/day
15	250	33.3
20	250	25.0
25	250	20.0
50	250	10.0