

**Neomycin Sulfate and Oxytetracycline Chicken Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated chicken feed.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....10 grams/lb.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 10 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee



INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below.

Indications for Use	Oxytetracycline and Neomycin Sulfate Amount	Mixing Directions
For control of infectious synovitis caused by <u>Mycoplasma synoviae</u> ; control of fowl cholera caused by <u>Pasteurella multocida</u> susceptible to oxytetracycline.	100 -200 g/ton of neomycin and oxytetracycline* *The final printed feed label should list only a single drug concentration	Mix 10 to 20 lb of this Type B medicated feed with 1990 to 1980 lb of non-medicated feed to make one ton of Type C medicated feed containing 100-200 g/ton of neomycin and oxytetracycline
For control of chronic respiratory disease (CRD) and air sac infection caused by	400 g/ton of neomycin and oxytetracycline	Mix 40 lb of this Type B medicated feed with 1960 lb of non-medicated feed to make one

<u>Mycoplasma gallisepticum</u> and <u>Escherichia coli</u> susceptible to oxytetracycline.		ton of Type C medicated feed containing 400 g/ton of neomycin and oxytetracycline
For reduction of mortality due to air sacculitis (air-sac-infection) caused by <u>Escherichia coli</u> susceptible to oxytetracycline.	500 g/ton of neomycin and oxytetracycline	Mix 50 lb of this Type B medicated feed with 1950 lb of non-medicated feed to make one ton of Type C medicated feed containing 500 g/ton of neomycin and oxytetracycline
<p style="text-align: center;">  Residue Warning: Do not feed to chickens producing eggs for human consumption. At 500 g/ton level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower uses. In low calcium feeds, withdraw 3 days before slaughter.  </p>		

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000