Neomycin Sulfate and Oxytetracycline Swine Ration B Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For treatment of bacterial enteritis caused by <u>Escherichia coli</u> and <u>Salmonella choleraesuis</u> and bacterial pneumonia caused by <u>Pasteurella multocida</u> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <u>Escherichia coli</u> susceptible to neomycin.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate	grams/ton.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to	<u></u> &
See FEEDING DIRECTIONS.	

GUARANTEED ANALYSIS

Crude Protein, (Min)	XX.XX%
Lysine, (Min)	X.XX%
Methionine, (Min)	X.XX%
Crude Fat, (Min)	X.XX%
Crude Fiber, (Max)	X.XX%
Calcium, (Min)	X.XX%
Calcium, (Max)	X.XX%
Phosphorus, (Min)	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Selenium, (Min)	ppm
Zinc, (Min)	ppm

¹If added

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 10mg/lb body weight/day continuously for 7-14 days as the sole ration. For example, the table below with the stated drug concentration, body weight and feed consumption will provide swine with 10 mg/lb body weight/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C feed	Body Weight (lb)	Type C Feed Consumption (lb)
400	100	5
500	150	6.00
600	200	6.67

RESIDUE WARNIN

RESIDUE WARNING: Withdraw 5 days before slaughter.



*The final printed medicated feed label must state a single drug concentration.

Net Weight____lb. (____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee