Chlortetracycline and Sulfamethazine Swine Ration

TYPE B MEDICATED FEED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

FOR USE IN THE MANUFACTURE OF SWINE FEEDS

| Active Drug Ingredients |
|--|
| Chlortetracycline calcium complex |
| Equivalent to chlortetracycline HCL 10 g/lb |
| Sulfamethazine2.2% (10 g/lb) |
| Indications: For reduction of the incidence of cervical abscesses, treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i> and vibrionic dysentery); prevention of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i> and vibrionic dysentery) during time of stress; maintenance of weight gains in the presence of atrophic rhinitis. |
| Guaranteed Analysis |
| Crude Protein, (Min)XX.XXX |
| Lysine, (Min)X.XX% |
| Crude Fat, (Min) |
| Crude Fiber, (Max)X.XX% |
| Calcium, (Min) |
| Calcium, (Max) |
| Phosphorus, (Min)X.XX% |
| Salt, (Min) ¹ |
| Salt, (Max) ¹ X.XX% |
| Total Sodium, (Min) ² X.XX% |
| Total Sodium, (Max) ² X.XX% |
| Selenium, (Min)ppm |
| Zinc, (Min)ppm |
| ¹ If added |
| ² Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee |
| Mixing Directions: Mix at the rate of 10 pounds (4.54 kg) per ton (907.2 kg) with part of the feed |
| ingredients to make a preblend. Add the remainder of the ingredients to give a final concentration of |
| 100 g chlortetracycline and 100 g sulfamethazine per ton of feed and mix thoroughly. |
| WARNING: WITHDRAW 15 DAYS PRIOR TO SLAUGHTER. |
| Lot No.: |
| Manufactured by |

BLUE BIRD FEED COMPANY BLUE BIRD, MD 00000

| Net Weight | : Lb | s. (| Kg. |
|------------|------|------|-----|
|------------|------|------|-----|