Chlortetracycline Chicken Ration II Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of chronic respiratory disease (CRD) and air sac infection caused by <u>Mycoplasma</u> gallisepticum and <u>Escherichia coli</u> susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Crude Protein, not less than	%
Crude Fat, not less than	
Crude Fiber, not more than	
Lysine (min)	%
Methionine (min)	
Calcium (min)	%
Calcium (max)	
Phosphorus(min)	%
Salt ¹ (min)	%
Salt ¹ (max)	%
Sodium ² (min)	
Sodium ² (max)	
¹ If added	

INGREDIENTS

²Should be guaranteed only when total sodium exceeds that furnished by

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7 to 14 days as the sole ration.

WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.

the maximum salt guarantee

*The final printed label must bear a single drug concentration.

Net Weight____lb. (____kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000