Chlortetracycline Turkey Ration IV Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline to provide 25 mg/lb body weight/day varying with body weight and feed consumption.

GUARANTEED ANALYSIS

Crude Protein, not less than	%
Crude Fat, not less than	%
Crude Fiber, not more than	%
Lysine (min)	%
Methionine (min)	
Calcium (min)	
Calcium (max)	
Phosphorus(min)	
Salt ¹ (min)	
Salt ¹ (max)	
Sodium ² (min)	
Sodium ² (max)	
¹ If added	

¹If added

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

For example, the table below with the stated drug concentration, body weight and feed consumption will provide turkeys with 25 mg/lb body weight/day of chlortetracycline.

For Tom Turkeys:

Age	Body Weight	Feed Consumption	Feed Consumption	Required Chlortetracycline level
(week	<u>s)(lbs)</u>	(lbs. per week)	(lbs. per day)	(grams/ton)
1	0.3	0.2	0.03	500
5	3.6	1.9	.3	600
10	12.2	5.2	0.7	871
15	26.4	8.2	1.2	1100
20	36.6	10.4	1.5	1210
24	42.8	11.6	1.7	1347

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

For Hen Turkeys:

_	D 1 XX 1	F 10	F 10	D ! 1011 1! 1 1
Α	Age Body Weigh	t Feed Consumption	Feed Consumption	Required Chlortetracycline level
(1	weeks) (lbs)	(lbs. per week)	(lbs. per day)	(grams/ton)
1	0.3	0.2	0.03	500
5	3.1	1.4	0.2	775
1	0 9.7	3.7	0.5	970
1:	5 18.3	6.2	0.9	1017
2	0 25.4	7.5	1.1	1154

Feed continuously for 7 to 14 days as the sole ration.

WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.

Net Weight___lb (___kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000