DERA.5/BOV ROG— CATTLE FEED TYPE C Medicated Feed

(chlortetracycline and lasalocid Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For increased rate of weight gain and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in pasture cattle over 700 pounds (slaughter, stocker, feeder cattle, beef replacement heifers).

ACTIVE DRUG INGREDIENTS*

Chlortetracycline ^a	25 to 1100 g/ton
Lasalocid ^b (as lasalocid sodium)	.30 to 600 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than	%
(This includes not more than% equivalent crude protein from non-protein nitrogen) ¹	
Crude fat, not less than	%
Crude fiber, not more than	
Calcium, not less than	<u>%</u>
Calcium, not more than	%
Phosphorus, not less than	%
Potassium, not less than	%
Salt ¹ , not less than	
Salt ¹ , not more than	%
Sodium ² , not less than Sodium ² , not more than	%
Sodium ² , not more than	%
Vitamin A ¹ , not less than	
¹ If added	

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously on a hand-fed basis 0.5 mg chlortetracycline per lb. body weight per day and not less than 60 mg or more than 300 mg lasalocid per head daily in at least 1 pound of feed for pasture cattle (slaughter, stocker, feeder cattle, and beef replacement heifers). Daily lasalocid intakes in excess of 200 mg/head/day in pasture cattle have not been shown to be more effective than 200 mg lasalocid/head/day. The following table provides examples of feeding rates and amounts⁻

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt. lb	lb/head/day	Amount	Feeding Rate	Amount	Feeding Rate
	70 Douy Wi.	10/11eau/uay	mg/lb BW/day	g/ton	mg/head/day	g/ton
700	0.14	1	0.5	700	300	600
	0.5	3.5	0.5	200	200	114.3
	2	14	0.5	50	210	30
1000	0.2	2	0.5	500	150	150
	0.4	4	0.5	250	200	100

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.



Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.



MANUFACTURED BY

Blue Bird Feed Company Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

^aDeracin® is the proprietary name of chlortetracycline Type A medicated article (ANADA 200-510). ^bBOVATEC is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Pharmgate LLC [March 2017]