

BLUE BIRD LABELING
chlortetracycline hydrochloride
Type B Medicated Turkey Poultry Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For reduction of mortality due to paratyphoid caused by *Salmonella typhimurium* susceptible to chlortetracycline in turkey poults not over 4 weeks of age.

Active Drug Ingredient

Chlortetracycline* 401 – 80,000 g/ton**

Guaranteed Analysis

Crude Protein (Min) %
 Lysine (Min) %
 Methionine (Min) %
 Crude Fat (Min) %
 Crude Fiber (Max) %
 Calcium (Min) %
 Calcium (Max) %
 Phosphorus (Min) %
 Salt^a (Min) %
 Salt^a (Max) %
 Sodium^b (Min) %
 Sodium^b (Max) %

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.
^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkey poults containing 400 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	80	1920	400
40,000 (20 g/lb)	20	1980	400
80,000 (40 g/lb)	10	1990	400

The resulting Type C medicated feed should be fed continuously as the sole ration to turkey poults not over 4 weeks of age.

Warning

Do not feed to turkeys producing eggs for human consumption.

Manufactured By:

Blue Bird Feed Mill
 City, State, Zip

NET WEIGHT ON BAG OR BULK

*Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).

**The final printed feed label should list only a single drug concentration.