BLUE BIRD LABELING chlortetracycline hydrochloride Type B Medicated Turkey Feed – SYN

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of infectious synovitis caused by Mycoplasma synoviae susceptible to chlortetracycline in turkeys.

Active Drug Ingredient Chlortetracycline* 201 – 80,000 g/ton** Guaranteed Analysis Crude Protein (Min) _____% Lysine (Min) _____% Methionine (Min) _____% Crude Fat (Min) ______% Crude Fiber (Max) ______% Calcium (Min) _______% Calcium (Max) _______% Phosphorus (Min) ________% Salt³ (Min) _________%

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 200 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	40	1960	200
20,000 (10 g/lb)	20	1980	200
40,000 (20 g/lb)	10	1990	200
80,000 (40 g/lb)	5	1995	200

The resulting Type C medicated feed should be fed continuously as the sole ration for 7 to 14 days.

Warning

Do not feed to turkeys producing eggs for human consumption.

Manufactured By:

Blue Bird Feed Mill City, State, Zip

NET WEIGHT ON BAG OR BULK

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

^{*}Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).

^{**}The final printed feed label should list only a single drug concentration.