BLUE BIRD LABELING

Bacitracin methylenedisalicylate

Type B medicated broiler & replacement chickens/growing turkey/growing pheasant feed – ROG/FE (bacitracin Type B medicated feed)

Do Not Feed Undiluted

For increased rate of weight gain and improved feed efficiency in broiler and replacement chickens, growing turkeys, and growing pheasants.

Active Drug Ingredient

Guaranteed Analysis

Guaranteea Analysis	
Crude Protein (Min)	%
Lysine (Min)	%
Methionine (Min)	%
Crude Fat (Min)	%
Crude Fiber (Max)	%
Calcium (Min)	%
Calcium (Max)	%
Phosphorus (Min)	%
Salt ^a (Min)	%
Salt ^a (Max)	%
Sodium ^b (Min)	%
Sodium ^b (Max)	%

Ingredients

Feed ingredients, as defined by AAFCO, must be listed on the final printed Type C medicated feed label by their common or usual names in descending order or predominance by weight.

Mixing Directions

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for [broiler chickens, replacement chickens, growing turkeys, growing pheasants] containing 4 – 50 g bacitracin per ton.

The following table provides examples of mixing rates:

Type B bacitracin concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C bacitracin concentration (g/ton)
10,000 (5 g/lb)	5	1995	25
10,000 (5 g/lb)	10	1990	50
50,000 (25 g/lb)	2	1998	50

The resulting Type C medicated feed should be administered continuously throughout the feeding period.

Manufactured By:

Blue Bird Feed Mill City, State, Zip

NET WEIGHT ON BAG OR BULK

Lot #

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

^{*}Pennitracin MD is the proprietary name of bacitracin methylenedisalicylate formulation (Type A Medicated Article).

^{**}The final printed feed label should list only a single drug concentration.