

**Bacitracin methylenedisalicylate NE-P – Chicken Feed**  
(bacitracin methylenedisalicylate Type B medicated feed)

Do Not Feed Undiluted.

**INDICATIONS FOR USE**

For the prevention of mortality caused by necrotic enteritis associated with *Clostridium perfringens* in broiler and replacement chickens.

**ACTIVE DRUG INGREDIENTS**

Bacitracin (as feed grade bacitracin methylenedisalicylate)<sup>a</sup>..... 50,000 g/ton  
(25 g/lb)

**GUARANTEED ANALYSIS**

Crude protein, not less than	.....	_____	%
Lysine, not less than	.....	_____	%
Methionine, not less than	.....	_____	%
Crude fat, not less than	.....	_____	%
Crude fiber, not more than	.....	_____	%
Calcium, not less than	.....	_____	%
Calcium, not more than	.....	_____	%
Phosphorus, not less than	.....	_____	%
Salt <sup>1</sup> , not less than	.....	_____	%
Salt <sup>1</sup> , not more than	.....	_____	%
Sodium <sup>2</sup> , not less than	.....	_____	%
Sodium <sup>2</sup> , not more than	.....	_____	%

<sup>1</sup> If added

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**MIXING DIRECTIONS**

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed containing 50 g bacitracin per ton.

The following table provides an example of mixing rates:

Type B medicated feed bacitracin concentration (g/ton)	Type B medicated feed per ton of Type C medicated feed (lb)	Non-medicated feed per ton of Type C medicated feed (lb)	Type C medicated feed bacitracin concentration (g/ton)
50,000 (25 g/lb)	2	1998	50

The resulting Type C medicated feed should be fed as the sole ration for 28 to 35 days, starting from the time chicks are placed for brooding.



**WARNING**



**Withdrawal Periods:** No withdrawal period is required when used according to labeling.

**Questions/Comments?** Contact Pharmgate Animal Health LLC at 1-800-380-6099 or [www.pharmgate.com](http://www.pharmgate.com). To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 141-137

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

Lot Number (if applicable): \_\_\_\_\_

<sup>a</sup>Pennitracin MD 50G® is the proprietary name of the bacitracin methylenedisalicylate formulation (Type A medicated article) approved by FDA under NADA # 141-137.