

Bacitracin methylenedisalicylate NE-P – Chicken Feed
(bacitracin methylenedisalicylate Type C medicated feed)

INDICATIONS FOR USE

For the prevention of mortality caused by necrotic enteritis associated with *Clostridium perfringens* in broiler and replacement chickens.

ACTIVE DRUG INGREDIENTS

Bacitracin (as feed grade bacitracin methylenedisalicylate)^a.....50 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than	_____	%
Lysine, not less than	_____	%
Methionine, not less than	_____	%
Crude fat, not less than	_____	%
Crude fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹ If added

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed as the sole ration for 28 to 35 days, starting from the time chicks are placed for brooding.

WARNING

Withdrawal Periods: No withdrawal period is required when used according to labeling.

Questions/Comments? Contact Pharmgate Animal Health LLC at 1-800-380-6099 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 141-137

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

Lot Number (if applicable): _____

*Pennitracin MD 50G® is the proprietary name of the bacitracin methylenedisalicylate formulation (Type A medicated article) approved by FDA under NADA # 141-137.