

**CALVES**  
**(oxytetracycline Type B medicated feed)**

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For use in manufacturing medicated calf feed for use in calf milk replacers or starter feeds.

**Indications for Use:** For treatment of bacterial enteritis caused by *Escherichia coli* organisms susceptible to oxytetracycline.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline hydrochloride<sup>a</sup>..... 2-20 g/lb\*

**GUARANTEED ANALYSIS**

Crude Protein (Min) .....	_____ %
NPN <sup>1</sup> (Max).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium <sup>1</sup> (Min).....	_____ %
Calcium <sup>1</sup> (Max).....	_____ %
Phosphorus <sup>1</sup> (Min).....	_____ %
Salt <sup>1</sup> (Min).....	_____ %
Salt <sup>1</sup> (Max).....	_____ %
Sodium <sup>2</sup> (Min).....	_____ %
Sodium <sup>2</sup> (Max).....	_____ %
Potassium <sup>1</sup> (Min).....	_____ %
Vitamin A <sup>3</sup> (Min) .....	_____ IU/lb

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.  
<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.  
<sup>3</sup> Other than precursors of Vitamin A, if added.

**INGREDIENTS**

Ingredients as defined by AAFCO

**MIXING DIRECTIONS**

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in  
Type C medicated feed

For Calves: Mix an amount of Type B feed to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for treatment of bacterial enteritis caused by *Escherichia coli* organisms susceptible to oxytetracycline. Feed continuously for 7-14 days as the sole ration.

10 mg/lb body weight/day

**Withdrawal Periods and Residue Warnings:** 5-day withdrawal period at 10 mg/lb dosage. This drug product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Approved by FDA under NADA # 138-938  
Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000

Lot number (if applicable) \_\_\_\_\_

\*The final printed medicated feed label must state a single drug concentration.

\_\_\_\_\_

<sup>a</sup> Pennox<sup>®</sup> is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [July 12, 2022]