

SHEEP
(oxytetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated sheep feed.

Indications for Use: Treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline in sheep.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride^a.....2-20 g/lb*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____ %
NPN ¹ (Max).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min)	_____ %
Calcium (Max)	_____ %
Phosphorus (Min)	_____ %
Salt ¹ (Min)	_____ %
Salt ¹ (Max)	_____ %
Sodium ² (Min).....	_____ %
Sodium ² (Max)	_____ %
Copper ^{3,4} (Min).....	_____ ppm
Copper ^{3,4} (Max).....	_____ ppm
Selenium ³ (Min)	_____ ppm
Vitamin A ⁵ (Min)	_____ IU/lb

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

³ Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

⁴ Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.

⁵ Other than precursors of Vitamin A, if added.

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

For Sheep: Mix an amount of Type B feed to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline to be fed continuously for 7-14 days.

10 mg/lb body weight/day



Withdrawal Periods: 5-day withdrawal period.



Net Weight _____ lb (_____ kg)

Approved by FDA under NADA # 138-938

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

Lot number (if applicable) _____

*The final printed medicated feed label must state a single drug concentration.

^a Pennox[®] is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [July 21, 2022]