

**TURKEYS - HEXAMITIASIS**  
**(oxytetracycline Type C medicated feed)**

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**Indications for Use:** Control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline in turkeys.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride<sup>a</sup>.....100 g/ton

**GUARANTEED ANALYSIS**

Crude Protein (Min) .....	_____ %
Lysine (Min).....	_____ %
Methionine (Min) .....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min) .....	_____ %
Calcium (Max) .....	_____ %
Phosphorus (Min).....	_____ %
Salt <sup>1</sup> (Min).....	_____ %
Salt <sup>1</sup> (Max) .....	_____ %
Sodium <sup>2</sup> (Min).....	_____ %
Sodium <sup>2</sup> (Max) .....	_____ %

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO

**FEEDING DIRECTIONS**

Feed continuously for 7-14 days as the sole ration.



**Withdrawal Periods and Residue Warnings:** Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.



Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Approved by FDA under NADA # 138-938  
Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000

Lot number (if applicable) \_\_\_\_\_

<sup>a</sup> Pennox<sup>®</sup> is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [June 24, 2022]