

# Product Profile

## MycoGard<sup>®</sup> One-Time



### *Mycoplasma hyopneumoniae* Bacterin

#### Product Description

- One-dose *Mycoplasma hyopneumoniae* bacterin for vaccination of swine
- Inactivated bivalent-adjuvanted formulation using StimGard<sup>™</sup> adjuvant
- Each vial contains 50 or 250 doses (1 mL/dose)



#### Advantages

- 1-mL dose – easy dose size for nursing piglets.
- Convenient dosing – allowing for very early *Mycoplasma hyopneumoniae* vaccination.
- Safe – easy on animals and staff.
- DOI of 164 days
- Colored cap - for ease of identification.

#### Dosage / Administration

- Shake well before use. Using aseptic technique, inject a single 1.0-mL dose intramuscularly (IM).

#### Indications / Uses

- For use in healthy piglets of all breeds, two weeks of age (14 days) or older, against *Mycoplasma hyopneumoniae*.
- Visit USDA website for full label indications & technical information. ([productdata.aphis.usda.gov](http://productdata.aphis.usda.gov)).

#### Precautions

- CAUTION: For animal use only.
- Store at 2 to 8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening.
- Do not vaccinate within 21 days before slaughter.
- Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative.
- This product has not been tested in pregnant animals.

#### Packaging

- 50-mL or 250-mL vials.
- Other sizes available.

#### Ordering Information

- Prod. number:  
20010-50 (50 mL)  
20010-250 (250 mL)

## Key Features

- Excellent syringeability.
- Advanced purification process allows low-volume injection for minimal tissue reaction.
- Cap colored for easy identification.
- QR (quick response) codes on the label for inventory and tracking purposes.

The labeling contains complete use information, including any cautions and warnings. Always read, understand and follow the labeling and use directions. See the reverse side for use directions and additional information.



**Pharmgate**  
ANIMAL HEALTH

1800 Sir Tyler Drive  
Wilmington, NC | 28405  
910.679.8364  
[Pharmgate.com](http://Pharmgate.com)

# Mycogard® One-Time

*Mycoplasma hyopneumoniae* Bacterin

Mycoplasma  
Hyopneumoniae  
Bacterin

Mycogard®  
One-Time | 50 ML  
50 DOSES



This product has been shown to be effective for the vaccination of healthy piglets, two weeks of age or older against Mycoplasma Hyopneumoniae. The duration of immunity is at least 164 days. For more information regarding efficacy and safety data, see [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

**ADMINISTRATION AND DOSAGE:** Dose: 1.0 ml intramuscularly (I.M.).

**CAUTION:** For animal use only. Do not mix with other products. In case of human exposure, contact a physician. Store at 2 to 8°C (35°F to 45°F). Do not freeze. Shake well before use. Use entire contents upon opening. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. For advice on revaccination frequency, consult your veterinarian.

Serial No:

Exp. Date:

Manufactured by:

**Pharmgate Biologics Inc.**  
2635 University Avenue  
West, Suite 140,  
St. Paul, MN 55114  
612-256-0930

**VLN 329 / PCN 2775.01**

Visit USDA website for full label indications & technical information: [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov)

## Pharmgate Animal Health swine vaccines/bacterins.

Product	Doses	Pharmgate Product Number
Mycogard® (2 dose)	50	20020-50
Mycogard® (2 dose)	250	20020-250
Mycogard-1™ (1 dose)	50	21010-50
Mycogard-1™ (1 dose)	250	21010-250
CircoGard® (1 dose)	50	22010-50
CircoGard® (1 dose)	250	22010-250
Circo/Mycogard® (1 dose)	50	25010-50
Circo/Mycogard® (1 dose)	250	25010-250

