

Product Profile

MycoGard[®]

Mycoplasma hyopneumoniae Bacterin

Product Description

- Two-dose *Mycoplasma hyopneumoniae* bacterin for vaccination of swine
- Inactivated bivalent-adjuvanted formulation using StimGard[™] adjuvant
- Each vial contains 50 or 250 doses (1 mL/dose)



Advantages

- 1-mL dose – convenient dose size for nursing piglets.
- Convenient dosing – allowing for very early *M hyopneumoniae* vaccination.
- Safe – easy on animals and staff.
- StimGard[™] bivalent adjuvant – promotes a robust and balanced immune response.
- Colored cap - for ease of identification.

Dosage / Administration

- Shake well before use. Using aseptic technique, inject the first 1.0-mL dose intramuscularly (IM). Administer the second 1.0-mL dose IM two weeks after the first dose.

Indications / Uses

- For use in healthy piglets of all breeds, 2 weeks of age or older, as an aid in the reduction of lung lesions caused by *Mycoplasma hyopneumoniae*.

Precautions

- CAUTION: For animal use only.
- Store at 2 to 8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening.
- Do not vaccinate within 21 days before slaughter.
- Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative.

Packaging

- 50-mL or 250-mL vials.
- Other sizes available.

Ordering Information

- Prod. number:
20020-50 (50 mL)
20020-250 (250 mL)

Key Features

- Pig-proven under field conditions.
- Contains StimGard[™] bivalent adjuvant.
- Proven in millions of vaccinated pigs.
- Excellent syringeability.
- Advanced purification process allows low-volume injection for minimal tissue reaction.
- Cap colored for easy identification.
- QR (quick response) codes on the label for inventory and tracking purposes.

The labeling contains complete use information, including any cautions and warnings. Always read, understand and follow the labeling and use directions. See the reverse side for use directions and additional information.



Pharmgate
ANIMAL HEALTH

1800 Sir Tyler Drive
Wilmington, NC | 28405
910.679.8364
Pharmgate.com

MycoGard®

Mycoplasma hyopneumoniae Bacterin

Mycoplasma
Hyopneumoniae
Bacterin

MycoGard® | 250 ML
250 DOSES



This inactivated vaccine is fortified with a novel adjuvant and presented in the form of a bacterin. The product is for use in healthy piglets of all breeds, 2 weeks of age or older, by or under the supervision of a veterinarian, as an aid in the reduction of lung lesions due to swine enzootic pneumonia, caused by *Mycoplasma hyopneumoniae*.

Administration and Dosage: First Dose: 1.0 mL intramuscularly (I.M.), Second Dose: 1.0 mL I.M. must be administered two weeks after the first dose.

Caution: For animal use only. Do not mix with other products except as specified on this label. In case of human exposure, contact a physician. Store at 2-8°C (35-46°F). Do not freeze. Shake well before use. Use entire contents upon opening the container. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. Revaccination interval not established.

Product No:

Serial No:

Exp. Date:

Manufactured by:

Pharmgate Biologics
2575 University Avenue
West, Suite 100,
St. Paul, MN 55114
612-256-0930

VLN 329 / PCN 2775.00

Visit USDA website for full label indications & technical information: productdata.aphis.usda.gov

Pharmgate Animal Health swine vaccines/bacterins.

Product	Doses	Pharmgate Product Number
MycoGard® (2 dose)	50	20020-50
MycoGard® (2 dose)	250	20020-250
MycoGard-1™ (1 dose)	50	21010-50
MycoGard-1™ (1 dose)	250	21010-250
CircoGard® (1 dose)	50	22010-50
CircoGard® (1 dose)	250	22010-250
Circo/MycoGard® (1 dose)	50	25010-50
Circo/MycoGard® (1 dose)	250	25010-250