

## **Pennox** (oxytetracycline) Veterinary Feed Directive for use in Calves, Beef Cattle and Nonlactating Dairy Cattle

Client	
Busin Home Addre	
Phon	e #: Phone #:
Appro	ximate number of animals to be treated:
Locati	on of animals:
Specia	al Instructions and/or other animal identifications:
	tion, Drug Level in Medicated Feed, and Duration of Use (select one and specify the additional required information):
	A) For Growing Cattle (over 400 lbs): For the reduction of the incidence of liver abscesses. Drug level: g/ton in order to provide 75 mg / head / day
	Duration of use: days
	B) For Cattle: For the prevention and treatment of the early stages of the shipping fever complex.  Drug level: g/ton in order to provide 0.5 to 2.0 g / head / day  Duration of use: days (3 to 5 days)
	C) For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.  Drug level: g/ton in order to provide 10 mg//lb body weight / day  Duration of use: days (7 to 14 days)
	Duration of use days (7 to 14 days)
Caut	ion: Use of feed containing this Veterinary Feed Directive (VFD) drug in a manner other than as directed on the labeling (extra-label use) is not permitted.
	For use in Dry Feeds Only. Not for Use in Liquid feed Supplements.
	Residue Warnings: 5-day withdrawal period at 10 mg/lb dosage [Indication C].  Zero day withdrawal period [Indications A and B].
Comb	ination Use:  This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s in combination with any other animal drugs.
VFD Is	suance Date: VFD Expiration Date: Month/Day/Year (Not to exceed 6 months from issuance date)
Veterir	arian's signature:

Original – Veterinarian Copy – Supplier Copy – Client